

Why is mucormycosis more difficult to cure than more common mycoses?

A. Katragkou^{1,2}, T. J. Walsh^{2,3,4} and E. Roilides¹

1) Infectious Diseases Unit, 3rd Department of Pediatrics, Faculty of Medicine, Aristotle University School of Health Sciences, Hippokration Hospital, Thessaloniki, Greece, 2) Transplantation-Oncology Infectious Diseases Program, 3) Department of Pediatrics and 4) Department of Microbiology & Immunology, Weill Cornell Medical Center, New York, NY, USA

Abstract

Although considered to be a rare infection, mucormycosis (zygomycosis) has emerged as the second most common invasive mould infection. Despite the advent of newer antifungal agents, mortality rate of mucormycosis remains exceedingly high. Successful management of mucormycosis requires early diagnosis, reversal of underlying predisposing risk factors, surgical debridement and prompt administration of active antifungal agents. However, mucormycosis is not always amenable to cure. There are challenging obstacles that lead to difficulties in management of amphotericin B. These include unique host-based risk factors for mucormycosis, the fungus' resistance to innate host defences and distinctive features of its immunopathogenesis, such as extensive angioinvasion, increased virulence and use of chelators by the fungus as siderophores. In addition to these obstacles, the difficulties in early diagnosis, including nonspecific clinical manifestations, lack of serological methods, as well limitations of culture and molecular methods, lead to delay in initiation of antifungal therapy. Finally, the variability of susceptibility to amphotericin B and resistance to most other conventional antifungal agents leads to major limitations in successful treatment of this devastating infection.

Keywords: Amphotericin B, *Aspergillus fumigatus*, mucormycosis, *Rhizopus oryzae*, zygomycosis

Article published online: 27 November 2013

Clin Microbiol Infect 2014; **20** (Suppl. 6): 74–81

Corresponding author: E. Roilides, 3rd Department of Pediatrics, Hippokration Hospital, Konstantinoupolis 49, GR-546 42 Thessaloniki, Greece
E-mail: roilides@med.auth.gr

Introduction

Invasive fungal infections are important causes of morbidity and mortality especially in immunocompromised patients. While *Candida* spp., *Aspergillus* spp. and *Cryptococcus neoformans* are responsible for the preponderance of invasive fungal infections, other less common fungal pathogens account for a significant proportion of morbidity and mortality [1]. Among these pathogens with worldwide distribution and increasing frequency are Zygomycetes causing mucormycosis (previously called zygomycosis) [2]. Mucormycosis is caused by fungi of the order Mucorales and the family Mucoraceae [3]. *Rhizopus*, *Mucor*, and *Lichtheimia* (formerly *Absidia*) are the most common genera that cause mucormycosis, accounting for 70–80% of all cases, whereas *Cunninghamella*, *Apophysomyces*, *Saksena*, *Rhizomucor*, *Cokeromyces*, *Actinomucor* and *Syncephalastrum* are responsible for <1–5% of reported cases [4].

A feature that is characteristic for Mucorales among other filamentous fungi is that they cause disproportionately serious infections with exceedingly high morbidity and mortality in relatively immunocompetent individuals [5]. This may be particularly observed in patients with diabetes mellitus, neonates, as well as those with wounds caused by surgery, trauma and burns. Despite aggressive surgical intervention and intensive antifungal treatment, mortality of mucormycosis is high, ranging from 50–100% depending on the disease form [3,5,6]. By comparison, mortality rates for candidiasis and aspergillosis range from 20–50% and 35–45%, respectively [7–9].

Successful management of mucormycosis requires early diagnosis, reversal of underlying predisposing risk factors,

surgical debridement and prompt administration of amphotericin B [10]. However, despite these measures, mucormycosis is not always amenable to cure. We will discuss, in this article, the challenging obstacles that lead to difficulties in management of mucormycosis (Table 1).

Patient Population at Risk for Mucormycosis

Mucormycosis is an infection affecting all ages, from premature neonates to elderly people with varying types of underlying conditions [5,11–13]. While the incidence rates of invasive mucormycosis in these diverse patient groups have not been well studied, the current data demonstrate increased incidence in a wide range of specific immunocompromised patient groups [2,5,11].

In the most contemporary epidemiologic study of mucormycosis in Europe, the most common underlying diseases were haematological malignancies (44%) followed by diabetes mellitus (17%) and trauma (17%) [2]. Data from a global fungal infection registry that included patients from central Europe and Asia demonstrated that malignancy (63%), diabetes (17%) and solid organ transplantation (10%) predominated as underlying conditions for mucormycosis [11].

A comprehensive review of 929 reported cases showed that the most common underlying condition were diabetes (36%), followed by malignancy (17%), solid organ transplantation (7%), deferoxamine therapy (6%), injection drug use (5%) and bone marrow transplantation (5%). In addition, the patients' underlying conditions were shown to be related with the pattern of infection, in that, patients with malignancy had

more likely pulmonary infection; whereas, patients with diabetes mellitus had sino-orbital and rhinocerebral disease [5,14]. This relationship would be, in all probability, a reflection of the host's immune function along with other pathological-anatomical factors [5].

However, it has also been postulated that the differences in disease pattern may reflect the diversity between developed and developing countries from where the cases were reported [15]. Specifically, in developed countries, most patients with mucormycosis are those with diabetes and haematological malignancies [6], while in developing countries those with uncontrolled diabetes and trauma predominate [16,17]. Notably, there is a relatively large proportion (c. 14% in children and 19% in adults) with no apparent immunocompromising conditions at the time of infection [5,18]. Within France, traumatic wounds in immunocompetent and immunocompromised patients comprise a distinct population [14].

The differences in the population at risk for mucormycosis depicted by the current studies could be explained by the period effect, publication bias or the contribution of different investigational centres in each study. However, it is characteristic that while other filamentous fungi affect more classically defined immunocompromised hosts, such as those with malignancies under chemotherapy, transplant recipients or inherited immunodeficiencies, mucormycosis can cause serious infections in relatively immunocompetent hosts, such as those with diabetes mellitus, on deferoxamine therapy, injection drug users and with no apparent immune defect other than a traumatic injury [5,18]. Correspondingly, taking also into account the ubiquitous nature of *Mucorales*, defining a specific population at risk for mucormycosis in order to target it is a challenging problem as 'candidate-patients' represent a quite heterogeneous group.

TABLE 1. Possible reasons for difficulty to treat mucormycosis

Variable underlying conditions of mucormycosis
Haematological malignancies
Diabetes
Solid organ transplantation
Haematopoietic stem cell transplantation
Injection drug use
Neonates
Trauma
Burns
Normal host
Deferoxamine therapy
Distinctive features of immunopathogenesis
Extensive angioinvasion
Increased virulence (exerted by some species, see text)
Use of chelators by fungus as siderophores
Difficulties in diagnosis
Non-specific clinical manifestations
Lack of validated or standardized serological methods
Difficulties in culturing organism
Limited availability of validated and standardized molecular methods
Limitations of antifungal therapy
Late or no diagnosis antemortem
Elevated MICs of amphotericin B
Higher MICs of <i>Cunninghamella</i> to amphotericin B
Triazole-specific variability of susceptibility
Species-specific variability of susceptibility to triazoles

Distinctive Features of the Immunopathogenesis of Mucormycosis

Medically important members of the order *Mucorales* share many features with other filamentous fungi such as portals of the host for infection (airways and disrupted mucocutaneous barriers), the main lines of innate host defences (phagocytes, specific ligands in fungal spores such as pathogen-associated molecular patterns (PAMPs) and immune cells such as Toll-like receptors (TLRs)), as well as histopathological and clinical features [19,20]. However, *Rhizopus oryzae* and other selected *Mucorales* possess unique virulence characteristics and exert distinctive host–pathogen interactions compared to other fungi facilitating, thus, host evasion and disease progression [21].

There are several lines of *in vitro* evidence showing that *R. oryzae* and other members of the Mucorales have reduced susceptibility to innate host defence as compared to other more common fungi, such as *Aspergillus fumigatus* or *Candida albicans* [22,23]. Moreover, differential interspecies susceptibility patterns to host responses exist within the order Mucorales [24–26]. Namely, members of the genus *Rhizopus* suffer less hyphal damage and stimulate an impaired oxidative burst in human phagocytes as compared to *Lichtheimia (Absidia)* spp. [24], and *Cunninghamella bertholletiae* shows, *in vitro*, increased resistance to phagocyte-induced hyphal damage and, *in vivo*, increased virulence in an experimental neutropenic pulmonary mucormycosis model in comparison with *Rhizopus* spp. [25,26]. In agreement are the results of the *Drosophila melanogaster* host model that simulates important aspects of mucormycosis in humans. In contrast to other fungi, species within the order Mucorales rapidly infect and kill *D. melanogaster* wild-flies, and their pathogenicity is linked with impaired phagocytic cell activity and hyphal damage compared with those of *A. fumigatus* [27]. These experimental findings are collectively consistent with epidemiological data and clinical experience showing greater prevalence of *Rhizopus* spp. compared to *Lichtheimia corymbifera* in immunocompromised patients [3,28–30] and increased mortality in patients with *C. bertholletiae* infection [4,5].

While the exact mechanisms underlying such variable responses against Zygomycetes have not yet been elucidated, the increased virulence exerted by some species has been associated with the induction of a more pronounced proinflammatory response by selected species [25,31]. It was postulated that differences in cell-wall constituents and ligands may lead to variable recognition of fungal cell wall recognition patterns by TLR and dectin receptors with consequent downstream altered expression of certain stimulatory molecules like chemokines and cytokines [24,31]. Indeed, the *D. melanogaster* model demonstrated the importance of fungal recognition for infection development showing that Toll-deficient flies exhibit increased susceptibility to infections caused by the Mucorales [27]. Whole-genome expression profiling in wild-type flies after infection with Mucorales vs. *A. fumigatus* revealed that genes acting on pathogen recognition, immune defence, stress response, detoxification, steroid metabolism or tissue repair are differentially regulated by these two fungal pathogens [27].

One of the critical characteristics in mucormycosis pathogenicity is the extensive angioinvasion that results in vessel thrombosis and tissue necrosis [6,32]. This angioinvasion leads to haematogenous dissemination of the organism to target organs, while ischaemic necrosis of the infected tissue can prevent leucocyte and antifungal agent penetration to the foci

of infection [21]. *R. oryzae* was used as a model system in understanding the basis of fungal pathogenicity. Sequencing the genome of a pathogenic *R. oryzae* strain there was evidence that the entire genome had been duplicated and retained two copies of three extremely sophisticated systems involved in energy generation and utilization. This gene duplication has led to the development of gene families related to fungal virulence, fungal cell wall synthesis enzymes and signal transduction, which may contribute to the invasive nature of *R. oryzae* [33].

The seminal clinical observations that patients with diabetic ketoacidosis as well as patients receiving dialysis who are treated with iron chelator deferoxamine are characteristically susceptible to mucormycosis highlights the central role of host iron in the pathogenesis of mucormycosis [21,34]. In proof of principle *in vitro* studies, it was shown that *Rhizopus* spp. can accumulate 8- and 40-fold greater amounts of iron supplied by deferoxamine than can *A. fumigatus* and *C. albicans*, respectively [35]. Similarly, data from animal models showed that administration of deferoxamine or free ions reduced survival of animals infected with *Rhizopus* spp. but not *C. albicans* [36–38]. Deferoxamine *per se* is not the pathogenetic factor for infection, but *Rhizopus* spp. utilize deferoxamine as a siderophore to supply previously unavailable iron to the fungus [6]. However, not all Zygomycetes have the same susceptibility to iron chelators [39].

Among the classic enhancers of mucormycosis pathogenicity in humans is corticosteroid immunosuppressive therapy and diabetes mellitus, which seem to impair the ability of macrophages to prevent germination of sporangiospores [40,41]. Neutropenia encountered in patients with haematological malignancies and in the pre-engraftment stage of HSCT is a major contribution to the pathogenesis of mucormycosis [20,21,23–26]. Further contributing to the pathogenesis of mucormycosis is the presence of a receptor–ligand interaction between endothelial cell surfaces and *Rhizopus oryzae* [21]. More recently, it was found that exposure of Mucorales to voriconazole selectively enhanced their virulence [42]. Up to date, a number of potential virulence factors including mycotoxins and lytic enzymes excreted by *Rhizopus* spp. have been proposed; however, their contribution to Zygomycetes pathogenicity needs to be further elucidated [21].

Difficulties in Mucormycosis Diagnosis

Early and accurate diagnosis is, *a priori*, the most critical aspect for improved outcome of mucormycosis given the limited therapeutic options available, which frequently involve disfiguring and debilitating surgeries. However, many suspected mucormycosis cases ranging from 4 to >90% are not

confirmed until post-mortem examination [2,43–45]. The establishment of a definite diagnosis is hampered by a variety of factors including non-specific clinical presentation of mucormycosis as well as the various drawbacks of the currently implemented diagnostic means.

There are some clinical manifestations such as diplopia, necrotic naso-sinus eschars, pleuritic pain, necrotic cutaneous lesions that, in the predisposed host, carry a potentially high predictive value [15,43]. These clinical manifestations are discussed in greater depth elsewhere. Nevertheless, these manifestations are nonspecific and their differential diagnosis includes a range of infections caused by angioinvasive pathogens, including *Aspergillus* spp., *Fusarium* spp., *Scedosporium* spp. and *Pseudomonas aeruginosa*; therefore, a high index of suspicion and prompt identification of host predisposing factors are required for early recognition of mucormycosis.

Cultural isolation and identification of the fungus to the genus or species level is of epidemiological, prognostic and therapeutic importance [4,26,43]. The cultural isolation yield ranges from 50 to 71%, while it was documented that it was significantly improved over time [2,5]. However, recovery of Mucorales from specimens in clinical microbiology laboratories is challenging. Performing invasive procedures to obtain the necessary material especially in severely ill patients with thrombocytopenia or coagulopathy may be a daunting and perilous task. The hyphae of Mucorales may be difficult to observe on wet mounts and need special chitin-binding stains with a fluorescent microscope or may not be abundant so as to be seen. Moreover, tissue handling aggressive tissue grinding or homogenization may destroy the coenocytic hyphae [43,46]. While Mucorales are usually morphologically distinctive from other filamentous fungi, in some cases where antifungal therapy has commenced before biopsy, morphological features may be atypical, reducing the ability to differentiate Mucorales from other filamentous fungi [43].

Currently, there are no readily available nonculture methods, such as measurement of biochemical or serological biomarkers, to facilitate the diagnosis of invasive mucormycosis. By comparison, circulating mannan antigen and (1→3)- β -D-glucan are used for diagnosis of invasive candidiasis, while galactomannan measured in serum and bronchoalveolar lavage (BAL) fluid is useful in the diagnosis of invasive aspergillosis.

Molecular identification is feasible and holds promise for early and accurate diagnosis of mucormycosis. Most studies have evaluated ribosomal targets (18S, 28S and internal transcribed spacer (ITS)) or other DNA targets (the high-affinity iron permease I gene *FTR1* or cytochrome b) that allow species identification of Zygomycetes from cultures [47]. Molecular methods also play a role in formalin-fixed, paraf-

fin-embedded biopsy samples, which, on many occasions, constitute the only material available for use for diagnosis in the clinical setting.

The greater challenge for early diagnosis of mucormycosis is to develop molecular systems for identification of amplicon from blood or BAL fluid in patients with invasive pulmonary mucormycosis. One of the first laboratory animal studies to demonstrate circulating species-specific amplicon from plasma and BAL fluid in experimental invasive pulmonary mucormycosis found sensitivities of 67% and 100% with a high degree of specificity [48]. Other studies from experimental animal models as well as case reports from patients further demonstrate the feasibility in identifying Mucorales such as *C. bertholletiae*, *Rhizomucor pusillus*, *R. microsporus*, *L. corymbifera* and *Saksenaea vasiformis* from tissue and cultures [48–55]. In these cases molecular approaches are mainly based on PCR assays; however, the performance of these techniques can become compromised, as formalin fixation is associated with DNA damage. Currently, molecular diagnostics for mucormycosis is not widely available; its use is compromised by limited sensitivity, time to detection and ability to provide rapid results. Collectively, while molecular diagnosis of mucormycosis is feasible, several areas, such as DNA extraction methodology, identification of informative DNA targets, a validated sequence database, a broader spectrum of primers, as well as systematic validation and standardization of assays in human specimens warrant improvement [43,56].

A protocol of the European Confederation of Medical Mycology (ECMM)—International Society for Human and Animal Mycology (ISHAM) Zygomycosis Working Group will be systematically addressing these needs for advancing the development of antigenic, biochemical and molecular assays for diagnosis of invasive pulmonary mucormycosis. The three objectives of development of a predictive risk model, an archive of human specimens and an educational effort are aimed at advancing the early diagnosis of invasive mucormycosis.

Diagnostic imaging and especially CT scan is an invaluable tool to aid in early diagnosis of invasive infections in immunocompromised hosts. Nevertheless, radiological features are not pathognomonic of mucormycosis while their absence cannot exclude this infection [57]. Many lesions, such as nodules, halo sign, reverse halo sign, cavities, wedge-shaped and pleural effusions, characteristically occur in pulmonary mucormycosis as well as other angioinvasive organisms like *Aspergillus* spp., *Scedosporium* spp., *Fusarium* spp. and *P. aeruginosa* [43,57].

Early and accurate diagnosis of mucormycosis is an area of critical importance and until newer molecular diagnostic techniques and biomarkers become available, diagnosis will

be based on prompt recognition of risk factors, clinical manifestations and radiological findings and with confirmation by culture and biopsy [43].

Difficulties in Treatment of Mucormycosis

The advent of newer antifungal agents significantly improved the prognosis in immunocompromised patients with invasive fungal infections. However, in the case of mucormycosis, the invasive nature of the disease leads to an overall mortality exceeding 50%, while on many occasions, antifungal therapy alone is rarely effective resulting in 100% mortality particularly for patients with disseminated disease [2,5,14] (Fig. 1).

Various Mucorales organisms have differential responses to antifungal agents. For example, *R. oryzae* tends to exhibit *in vitro* resistance to posaconazole; *Mucor circinelloides* shows greater susceptibility to posaconazole, and *Cunninghamella* tends to have higher MICs to amphotericin B [43]. In addition, the same species have variable response to the same antifungal class, for example *R. oryzae* has a variable response to different triazoles, including lack of activity of voriconazole as compared to relatively good activity of posaconazole. The recent sequencing of *R. oryzae* revealed that this strain is genetically equipped for adaptation to hostile environments such as the effects of antifungal agents. It has been advocated that the variable responses of *R. oryzae* to voriconazole, itraconazole and posaconazole are possibly due to the increased copy number and divergence of duplicated *ERG11* (the principal gene target for the triazoles) [33]. Consequently, the MICs for itraconazole and posaconazole are 4–8 dilutions higher for *R. oryzae* than those for *A. fumigatus*, and fungicidal activity is not achieved over a range of safely achievable drug concen-

trations [58–60]. The frequent breakthrough *R. oryzae* infections occurring in patients receiving voriconazole indicate that this antifungal agent is ineffective against *R. oryzae* [61,62]. This adaptive ability of *R. oryzae* may result in higher potential for development of resistance during long-term triazole therapy than *A. fumigatus* [63].

Amphotericin B is considered the first line therapy for mucormycosis. However, there are many lingering use problems to resolve, including the optimal dosage and timing for treatment initiation for each site of infection. Many clinicians who treat mucormycosis in an effort to control the infection use maximum tolerance doses of liposomal amphotericin B with a risk for nephrotoxicity. Studies from animal models showed that higher amphotericin B tissue concentration may be required for effective treatment of mucormycosis as compared to aspergillosis [64–66], while a phase II clinical trial of high-dose liposomal amphotericin B (10 mg/kg/day, IV) in mucormycosis treatment has been recently completed pending the efficacy results (<http://clinicaltrials.gov/show/NCT00467883>).

The impact of prompt amphotericin B-based treatment on the outcome of patients with mucormycosis has been determined by a retrospective study [67]. Chamilos *et al.* [67] demonstrated that delayed amphotericin B-based therapy resulted in a two-fold increase in mortality in patients with haematological malignancy and mucormycosis compared with early treatment (83% vs. 49%). Furthermore, delayed treatment of invasive mucormycosis was an independent predictor of poor outcome in multivariate analysis (odds ratio, 8 (95% confidence interval, 1.7–38.2); *p* 0.008). Given that Mucorales grow rapidly *in vivo*, there is a 'window of opportunity', which is much shorter than that of aspergillosis, where effective treatment should be initiated before extensive angioinvasion and dissemination occur [63]. Notably, while *in vitro* echinocandins demonstrate virtually no activity against Mucorales, *in vivo*, they are modestly effective [63]. Notably, at higher doses, echinocandins have attenuated activity compared with lower doses possibly reflecting upregulation of homeostatic cell-wall responses in the fungi that 'rescue' the fungus from the effects of echinocandins through compensatory increase in chitin synthesis [68,69]. In this regard, treatment options with predictable and favourable pharmacokinetics, such as lipid formulations of amphotericin B with echinocandin provide a reasonable rationale for phase III randomized, double-blinded, placebo-controlled trials [63]. However, more work on the safety, tolerability, efficacy, pharmacokinetics and pharmacodynamics is needed in a wider range of animal models of invasive pulmonary mucormycosis to assure that this hypothesis is tenable before embarking on a phase III trial.




FIG. 1. Treatment modalities against mucormycosis. PMN, polymorphonuclear neutrophils; MNC, monocytes; GM-CSF, Granulocyte-Macrophage-Colony-Stimulating Factor; IFN- γ , Interferon- γ .

Conclusions

Mucormycosis has a worse outcome than other invasive fungal infections such as candidiasis or aspergillosis. The higher degree of difficulty to cure this devastating infection is related to differences in host–fungus interactions, and pathogenetic mechanisms, as well as greater difficulties in early diagnosis when the ‘window’ of successful treatment is higher, and wider inadequacies of therapeutic options. Further advances in understanding host defence, developing newer diagnostic tools and creating better therapeutic interventions may improve outcome of this devastating disease.

Acknowledgements

Dr Walsh is a Scholar of the Henry Schueler Foundation in Mucormycosis and of the Sharp Family Foundation in Pediatric Infectious Diseases.

Transparency Declaration

Dr Katragkou does not disclose any potential conflict of interests. Dr Walsh receives research grants for experimental and clinical antimicrobial pharmacotherapeutics from Astellas, ContraFect, Trius, Cubist, Merck, Medimmune, Novartis and Pfizer. He has served as consultant to Astellas, ContraFect, Cubist, Drais, Gilead, iCo, Novartis, Methygene, Pfizer, SigmaTau and Trius. Dr Roilides has served on the speaker's bureau of Gilead, Cephalon, Pfizer, Wyeth, Schering, Merck, Aventis, Astellas, Elpen; has served as consultant to Schering, Gilead, Astellas, Pfizer; and has received grant support made to his institution from Pfizer, Gilead, Enzon, Schering, Wyeth.

References

- Walsh TJ, Groll A, Hiemenz J, Fleming R, Roilides E, Anaissie E. Infections due to emerging and uncommon medically important fungal pathogens. *Clin Microbiol Infect* 2004; 10 (suppl 1): 48–66.
- Skiada A, Pagano L, Groll A et al. Zygomycosis in Europe: analysis of 230 cases accrued by the registry of the European Confederation of Medical Mycology (ECMM) working group on zygomycosis between 2005 and 2007. *Clin Microbiol Infect* 2011; 17: 1859–1867.
- Ribes JA, Vanover-Sams CL, Baker DJ. Zygomycetes in human disease. *Clin Microbiol Rev* 2000; 13: 236–301.
- Gomes MZ, Lewis RE, Kontoyiannis DP. Mucormycosis caused by unusual mucormycetes, non-rhizopus, -mucor, and -lichtheimia species. *Clin Microbiol Rev* 2011; 24: 411–445.
- Roden MM, Zaoutis TE, Buchanan WL et al. Epidemiology and outcome of zygomycosis: a review of 929 reported cases. *Clin Infect Dis* 2005; 41: 634–653.
- Spellberg B, Edwards J Jr, Ibrahim A. Novel perspectives on mucormycosis: pathophysiology, presentation, and management. *Clin Microbiol Rev* 2005; 18: 556–569.
- Horn DL, Neofytos D, Anaissie EJ et al. Epidemiology and outcomes of candidemia in 2019 patients: data from the prospective antifungal therapy alliance registry. *Clin Infect Dis* 2009; 48: 1695–1703.
- Burgos A, Zaoutis TE, Dvorak CC et al. Pediatric invasive aspergillosis: a multicenter retrospective analysis of 139 contemporary cases. *Pediatrics* 2008; 121: e1286–e1294.
- Ruping MJ, Heinz WJ, Kindo AJ et al. Clinical epidemiology of 960 patients with invasive aspergillosis from the path alliance registry. *J Infect* 2012; 65: 453–464.
- Spellberg B, Walsh TJ, Kontoyiannis DP, Edwards J Jr, Ibrahim AS. Recent advances in the management of mucormycosis: from bench to bedside. *Clin Infect Dis* 2009; 48: 1743–1751.
- Ruping MJ, Heinz WJ, Kindo AJ et al. Forty-one recent cases of invasive zygomycosis from a global clinical registry. *J Antimicrob Chemother* 2010; 65: 296–302.
- Roilides E, Zaoutis TE, Walsh TJ. Invasive zygomycosis in neonates and children. *Clin Microbiol Infect* 2009; 15 (suppl 5): 50–54.
- Roilides E, Zaoutis TE, Katragkou A, Benjamin DK Jr, Walsh TJ. Zygomycosis in neonates: an uncommon but life-threatening infection. *Am J Perinatol* 2009; 26: 565–573.
- Lanternier F, Dannaoui E, Morizot G et al. A global analysis of mucormycosis in france: the retrozygo study (2005–2007). *Clin Infect Dis* 2012; 54 (suppl 1): S35–S43.
- Petrikkos G, Skiada A, Lortholary O, Roilides E, Walsh TJ, Kontoyiannis DP. Epidemiology and clinical manifestations of mucormycosis. *Clin Infect Dis* 2012; 54 (suppl 1): S23–S34.
- Prabhu RM, Patel R. Mucormycosis and entomophthoromycosis: a review of the clinical manifestations, diagnosis and treatment. *Clin Microbiol Infect* 2004; 10 (suppl 1): 31–47.
- Chakrabarti A, Das A, Mandal J et al. The rising trend of invasive zygomycosis in patients with uncontrolled diabetes mellitus. *Med Mycol* 2006; 44: 335–342.
- Zaoutis TE, Roilides E, Chiou CC et al. Zygomycosis in children: a systematic review and analysis of reported cases. *Pediatr Infect Dis J* 2007; 26: 723–727.
- Netea MG, Van der Meer JW, Kullberg BJ. Role of the dual interaction of fungal pathogens with pattern recognition receptors in the activation and modulation of host defence. *Clin Microbiol Infect* 2006; 12: 404–409.
- Roilides E, Kontoyiannis DP, Walsh TJ. Host defenses against zygomycetes. *Clin Infect Dis* 2012; 54 (suppl 1): S61–S66.
- Ibrahim AS, Spellberg B, Walsh TJ, Kontoyiannis DP. Pathogenesis of mucormycosis. *Clin Infect Dis* 2012; 54 (suppl 1): S16–S22.
- Chamilos G, Lewis RE, Lamas G, Walsh TJ, Kontoyiannis DP. Zygomycetes hyphae trigger an early, robust proinflammatory response in human polymorphonuclear neutrophils through toll-like receptor 2 induction but display relative resistance to oxidative damage. *Antimicrob Agents Chemother* 2008; 52: 722–724.
- Liles WC, Huang JE, van Burik JA, Bowden RA, Dale DC. Granulocyte colony-stimulating factor administered *in vivo* augments neutrophil-mediated activity against opportunistic fungal pathogens. *J Infect Dis* 1997; 175: 1012–1015.
- Gil-Lamaignere C, Simitsopoulou M, Roilides E, Maloukou A, Winn RM, Walsh TJ. Interferon gamma and granulocyte-macrophage colony-stimulating factor augment the activity of polymorphonuclear leukocytes against medically important zygomycetes. *J Infect Dis* 2005; 191: 1180–1187.
- Simitsopoulou M, Georgiadou E, Walsh TJ, Roilides E. *Cunninghamella bertholletiae* exhibits increased resistance to human neutrophils with or without antifungal agents as compared to *Rhizopus* spp. *Med Mycol* 2010; 48: 720–724.

26. Petraitis V, Petraitiene R, Antachopoulos C *et al.* Increased virulence of *Cunninghamella bertholletiae* in experimental pulmonary mucormycosis: correlation with circulating molecular biomarkers, sporangiospore germination and hyphal metabolism. *Med Mycol* 2013; 51: 72–82.
27. Chamilos G, Lewis RE, Hu J *et al.* *Drosophila melanogaster* as a model host to dissect the immunopathogenesis of zygomycosis. *Proc Natl Acad Sci USA* 2008; 105: 9367–9372.
28. Gonzalez CE, Rinaldi MG, Sugar AM. Zygomycosis. *Infect Dis Clin North Am* 2002; 16: 895–914, vi.
29. Kontoyiannis DP, Wessel VC, Bodey GP, Rolston KV. Zygomycosis in the 1990s in a tertiary-care cancer center. *Clin Infect Dis* 2000; 30: 851–856.
30. Marr KA, Carter RA, Crippa F, Wald A, Corey L. Epidemiology and outcome of mould infections in hematopoietic stem cell transplant recipients. *Clin Infect Dis* 2002; 34: 909–917.
31. Warris A, Netea MG, Verweij PE *et al.* Cytokine responses and regulation of interferon-gamma release by human mononuclear cells to *Aspergillus fumigatus* and other filamentous fungi. *Med Mycol* 2005; 43: 613–621.
32. Ben-Ami R, Luna M, Lewis RE, Walsh TJ, Kontoyiannis DP. A clinicopathological study of pulmonary mucormycosis in cancer patients: extensive angioinvasion but limited inflammatory response. *J Infect* 2009; 59: 134–138.
33. Ma LJ, Ibrahim AS, Skory C *et al.* Genomic analysis of the basal lineage fungus *Rhizopus oryzae* reveals a whole-genome duplication. *PLoS Genet* 2009; 5: e1000549.
34. Boelaert JR, van Roost GF, Vergauwe PL, Verbanck JJ, de Vroey C, Segaut MF. The role of desferrioxamine in dialysis-associated mucormycosis: report of three cases and review of the literature. *Clin Nephrol* 1988; 29: 261–266.
35. Boelaert JR, de Locht M, Van Cutsem J *et al.* Mucormycosis during deferoxamine therapy is a siderophore-mediated infection. *In vitro* and *in vivo* animal studies. *J Clin Invest* 1993; 91: 1979–1986.
36. Boelaert JR, Van Cutsem J, de Locht M, Schneider YJ, Crichton RR. Deferoxamine augments growth and pathogenicity of *Rhizopus*, while hydroxypyridinone chelators have no effect. *Kidney Int* 1994; 45: 667–671.
37. de Locht M, Boelaert JR, Schneider YJ. Iron uptake from ferrioxamine and from ferrirrhizoferrin by germinating spores of *Rhizopus microsporus*. *Biochem Pharmacol* 1994; 47: 1843–1850.
38. Van Cutsem J, Boelaert JR. Effects of deferoxamine, ferroxamine and iron on experimental mucormycosis (zygomycosis). *Kidney Int* 1989; 36: 1061–1068.
39. Lewis RE, Pongas GN, Albert N, Ben-Ami R, Walsh TJ, Kontoyiannis DP. Activity of deferasirox in mucorales: influences of species and exogenous iron. *Antimicrob Agents Chemother* 2011; 55: 411–413.
40. Shoham S, Levitz SM. The immune response to fungal infections. *Br J Haematol* 2005; 129: 569–582.
41. Waldorf AR, Levitz SM, Diamond RD. *In vivo* bronchoalveolar macrophage defense against *Rhizopus oryzae* and *Aspergillus fumigatus*. *J Infect Dis* 1984; 150: 752–760.
42. Lamarin GA, Ben-Ami R, Lewis RE, Chamilos G, Samonis G, Kontoyiannis DP. Increased virulence of Zygomycetes organisms following exposure to voriconazole: a study involving fly and murine models of zygomycosis. *J Infect Dis* 2009; 199: 1399–1406.
43. Walsh TJ, Gamaletsou MN, McGinnis MR, Hayden RT, Kontoyiannis DP. Early clinical and laboratory diagnosis of invasive pulmonary, extrapulmonary, and disseminated mucormycosis (zygomycosis). *Clin Infect Dis* 2012; 54 (suppl 1): S55–S60.
44. Pagano L, Offidani M, Fianchi L *et al.* Mucormycosis in hematologic patients. *Haematologica* 2004; 89: 207–214.
45. Ingram CW, Sennesh J, Cooper JN, Perfect JR. Disseminated zygomycosis: report of four cases and review. *Rev Infect Dis* 1989; 11: 741–754.
46. Monheit JE, Cowan DF, Moore DG. Rapid detection of fungi in tissues using calcofluor white and fluorescence microscopy. *Arch Pathol Lab Med* 1984; 108: 616–618.
47. Dannaoui E. Molecular tools for identification of zygomycetes and the diagnosis of zygomycosis. *Clin Microbiol Infect* 2009; 15 (suppl 5): 66–70.
48. Kasai M, Harrington SM, Francesconi A *et al.* Detection of a molecular biomarker for zygomycetes by quantitative PCR assays of plasma, bronchoalveolar lavage, and lung tissue in a rabbit model of experimental pulmonary zygomycosis. *J Clin Microbiol* 2008; 46: 3690–3702.
49. Schwarz P, Bretagne S, Gantier JC *et al.* Molecular identification of zygomycetes from culture and experimentally infected tissues. *J Clin Microbiol* 2006; 44: 340–349.
50. Machouart M, Larche J, Burton K *et al.* Genetic identification of the main opportunistic mucorales by pcr-restriction fragment length polymorphism. *J Clin Microbiol* 2006; 44: 805–810.
51. Iwen PC, Freifeld AG, Sigler L, Tarantolo SR. Molecular identification of *Rhizomucor pusillus* as a cause of sinus-orbital zygomycosis in a patient with acute myelogenous leukemia. *J Clin Microbiol* 2005; 43: 5819–5821.
52. Lechevalier P, Hermoso DG, Carol A *et al.* Molecular diagnosis of *Saksenaia vasiformis* cutaneous infection after scorpion sting in an immunocompetent adolescent. *J Clin Microbiol* 2008; 46: 3169–3172.
53. Kobayashi M, Togitani K, Machida H, Uemura Y, Ohtsuki Y, Taguchi H. Molecular polymerase chain reaction diagnosis of pulmonary mucormycosis caused by *Cunninghamella bertholletiae*. *Respirology* 2004; 9: 397–401.
54. Larche J, Machouart M, Burton K *et al.* Diagnosis of cutaneous mucormycosis due to *Rhizopus microsporus* by an innovative PCR-restriction fragment-length polymorphism method. *Clin Infect Dis* 2005; 41: 1362–1365.
55. Dannaoui E, Schwarz P, Slany M *et al.* Molecular detection and identification of zygomycetes species from paraffin-embedded tissues in a murine model of disseminated zygomycosis: a collaborative european society of clinical microbiology and infectious diseases (escmid) fungal infection study group (efisg) evaluation. *J Clin Microbiol* 2010; 48: 2043–2046.
56. Balajee SA, Sigler L, Brandt ME. DNA and the classical way: identification of medically important molds in the 21st century. *Med Mycol* 2007; 45: 475–490.
57. Georgiadou SP, Sipsas NV, Marom EM, Kontoyiannis DP. The diagnostic value of halo and reversed halo signs for invasive mold infections in compromised hosts. *Clin Infect Dis* 2011; 52: 1144–1155.
58. Sabatelli F, Patel R, Mann PA *et al.* *In vitro* activities of posaconazole, fluconazole, itraconazole, voriconazole, and amphotericin B against a large collection of clinically important molds and yeasts. *Antimicrob Agents Chemother* 2006; 50: 2009–2015.
59. Sun QN, Fothergill AW, McCarthy DI, Rinaldi MG, Graybill JR. *In vitro* activities of posaconazole, itraconazole, voriconazole, amphotericin b, and fluconazole against 37 clinical isolates of zygomycetes. *Antimicrob Agents Chemother* 2002; 46: 1581–1582.
60. Dannaoui E, Meletiadis J, Mouton JW, Meis JF, Verweij PE. *In vitro* susceptibilities of zygomycetes to conventional and new antifungals. *J Antimicrob Chemother* 2003; 51: 45–52.
61. Kontoyiannis DP, Lionakis MS, Lewis RE *et al.* Zygomycosis in a tertiary-care cancer center in the era of aspergillus-active antifungal therapy: a case-control observational study of 27 recent cases. *J Infect Dis* 2005; 191: 1350–1360.
62. Singh N, Aguado JM, Bonatti H *et al.* Zygomycosis in solid organ transplant recipients: a prospective, matched case-control study to assess risks for disease and outcome. *J Infect Dis* 2009; 200: 1002–1011.
63. Lewis RE, Lortholary O, Spellberg B, Roilides E, Kontoyiannis DP, Walsh TJ. How does antifungal pharmacology differ for mucormycosis versus aspergillosis? *Clin Infect Dis* 2012; 54 (suppl 1): S67–S72.

64. Ibrahim AS, Gebremariam T, Husseiny MI et al. Comparison of lipid amphotericin B preparations in treating murine zygomycosis. *Antimicrob Agents Chemother* 2008; 52: 1573–1576.
65. Lewis RE, Liao G, Hou J, Chamilos G, Prince RA, Kontoyiannis DP. Comparative analysis of amphotericin B lipid complex and liposomal amphotericin B kinetics of lung accumulation and fungal clearance in a murine model of acute invasive pulmonary aspergillosis. *Antimicrob Agents Chemother* 2007; 51: 1253–1258.
66. Lewis RE, Albert ND, Liao G, Hou J, Prince RA, Kontoyiannis DP. Comparative pharmacodynamics of amphotericin B lipid complex and liposomal amphotericin B in a murine model of pulmonary mucormycosis. *Antimicrob Agents Chemother* 2010; 54: 1298–1304.
67. Chamilos G, Lewis RE, Kontoyiannis DP. Delaying amphotericin b-based frontline therapy significantly increases mortality among patients with hematologic malignancy who have zygomycosis. *Clin Infect Dis* 2008; 47: 503–509.
68. Ibrahim AS, Bowman JC, Avanesian V et al. Caspofungin inhibits *Rhizopus oryzae* 1,3-beta-d-glucan synthase, lowers burden in brain measured by quantitative PCR, and improves survival at a low but not a high dose during murine disseminated zygomycosis. *Antimicrob Agents Chemother* 2005; 49: 721–727.
69. Walker LA, Munro CA, de Bruijn I, Lenardon MD, McKinnon A, Gow NA. Stimulation of chitin synthesis rescues *Candida albicans* from echinocandins. *PLoS Pathog* 2008; 4: e1000040.